Index

Note: Page numbers of article titles are in **boldface** type.

A

- Acute pain management service, setting up an, **893–910**
 - continuous peripheral nerve blocks, 903–907
 - monitoring for falls in patients with, 906–907
 - potential complications of, 903–906
 - role in ambulatory continuous perineural catheter service, 907
 - role of local anesthetic additives in, 907
- intravenous patient-controlled analgesia, 896–898
- multimodal approach to pain management, 894
- neuraxial analgesia, 900–903
- opioid-tolerant patients, 898–900
- role of nurses in and use of protocols, 896
- structure and individual roles within, 894–896
- Ambulatory continuous perineural catheter service, postoperative considerations for, 918
 - role of acute pain management service in, 907
- Ambulatory regional anesthesia program, for orthopedic surgery, **911–921**
 - postoperative considerations for ambulatory perineural catheters, 918
 - potential advantages of regional anesthesia, 911–913
 - decreases postoperative nausea and vomiting, 912
 - improves patient satisfaction, 913
 - increases efficiency, 912
 - optimizes analgesia, 911–912
 - prevents unanticipated admissions and leads to earlier discharge, 912
 - surgeons preference for, 913
 - practical details for, 913–918
 - equipment and personnel, 914
 - models of practice, 915–916
 - multimodal approach and clinical pathways, 916
 - patient education and screening, 913
 - postoperative conditions, 916–918
 - procedures amenable to perineural blockade, 915
 - techniques for block placement, 914–915
- Analgesia. *See* Pain management.
- Anesthesia, local. *See* Local anesthetics.
 - regional. *See* Regional anesthesia.
- Anesthesiologists, role in orthopedic trauma, 789–780
- Ankylosing spondylitis, preoperative evaluation for orthopedic surgery in patients with, 884–885

B

- Battlefield anesthesia, for orthopedic trauma, 790–791
- Blood conservation strategies, for orthopedic surgery, 888

Anesthesiology Clin 32 (2014) 933–941

[http://dx.doi.org/10.1016/S1932-2275\(14\)00122-0](http://dx.doi.org/10.1016/S1932-2275(14)00122-0)

anesthesiology.theclinics.com

1932-2275/14/\$ – see front matter © 2014 Elsevier Inc. All rights reserved.

C

- Cancer recurrence, relationship of regional anesthesia-analgesia to, **841–851**
 - immune system and cancer, 842
 - meta-analysis, 847
 - opioids and immune system, 843
 - regional and local anesthetic effect on immunosuppression, 843
 - studies investigating, 844–847
 - surgical site infections and regional anesthesia, 847–848
 - surgical stress leading to immunosuppression, 842–843
- Catheters, ambulatory continuous perineural, postoperative considerations for, 918
 - role of acute pain management service in, 907
- Central neuraxial anesthesia, evidence for ultrasound guidance in, 774–778
- Comfort, patient, with regional anesthesia for orthopedic surgery, 813–814
- Compartment syndrome, 800–802
 - risk with regional anesthesia for orthopedic trauma, 798–800
- Complications, with regional anesthesia for orthopedic surgery, 811–812
- Continuous perineural catheter service, role of acute pain management service in, 907
- Continuous peripheral nerve blocks, evidence for ultrasound guidance and, 779–780
 - in an acute pain service, 903–907
 - monitoring for falls in patients with, 906–907
 - potential complications of, 903–906
 - role in ambulatory continuous perineural catheter service, 907
 - role of local anesthetic additives in, 907
- Critical care admission, with regional anesthesia for orthopedic surgery, 812–813
- Crush injuries, in orthopedic trauma, regional techniques for, 793–794

D

- Discharge, earlier, with regional anesthesia for orthopedic surgery, 815

E

- Economic advantages, with regional anesthesia for orthopedic surgery, 814–815
- Education, patient, prior to orthopedic surgery, 888–889
 - in ambulatory regional anesthesia program, 913

F

- Fall reduction program, for lower-extremity joint arthroplasty patients, **853–864**
 - assessment, 856
 - current guidelines, 854–855
 - example of selective interventions for, 858–861
 - alert system, 860–861
 - bed-exit alarm, 861
 - education, 859–860
 - fall risk assessment tools, 860
 - preliminary results, 861
 - implementation, 858
 - multidisciplinary approach, 855
 - risk factors, 856–857

- strategies and interventions, 857
- sustainability, 858

Fall risk, inpatient, with regional anesthesia for orthopedic surgery, 815–816

- monitoring for, in patients with lower extremity blockade, 906–907

Functional outcome, with regional anesthesia for orthopedic surgery, 815–816

H

Hemophilia, preoperative evaluation for orthopedic surgery in patients with, 886–887

Hemostasis, in orthopedic trauma, 804

Hip fractures, optimizing perioperative care for patients with, **823–839**

- challenges of, 823–824

- institutional guidelines, 833

- intraoperative considerations, 831–833

- choice of anesthesia, 833

- surgical, 831–833

- perioperative risk and optimization, 824–831

- cardiac considerations, 824–826

- delirium, 829

- other medical considerations, 829–830

- pulmonary considerations, 826–829

- recommendation, 830

- risk of venous thromboembolism, 829

- postoperative care, 833

- preoperative evaluation and patient preparation for patients with, 883–884

I

Immune system, cancer and, 842

- opioids and, 843

Immunosuppression, regional anesthesia and local anesthetic effects on, 843

- surgical stress leading to, 842–843

Infection, risk with regional anesthesia for orthopedic trauma, 800–802

- surgical site, and regional anesthesia, 847–848

J

Joint replacement, developing a fall reduction program for patients undergoing total, **853–864**

K

Ketamine, for postoperative pain in opioid-tolerant patients, 927–929

L

Length of stay, with regional anesthesia for orthopedic surgery, 815

Local anesthetics, effects on immunosuppression, 843

- evidence for optimal ultrasound-guided distribution of, 780–782

- for regional anesthesia in orthopedic trauma, 797–798

- neuraxial blocks, 797–798

Local (*continued*)

- peripheral nerve blocks, 798
 - role in perineural analgesia, 907
- Lower-extremity joint arthroplasty, developing a fall reduction program for patients undergoing, **853–864**

M

- Medications, perioperative management in patients undergoing orthopedic surgery, 887–888
- Morbidity, with regional anesthesia for orthopedic surgery, 811–812
- Mortality, with regional anesthesia for orthopedic surgery, 811–812

N

- Nausea, postoperative, decrease in with regional anesthesia, 912
- Neuraxial techniques, in acute pain service, 900–903
 - in orthopedic trauma, 793
- Nociception, neurochemistry of, in trauma, 791–792
- Nurses, role in acute pain management service, 896

O

- Opioid-tolerant patients, in acute pain management service, 898–900
 - chronic users, 898–899
 - patients with opioid abuse or addiction, 899–900
- perioperative management for orthopedic surgery in, **923–932**
 - background on clinical studies, 926–927
 - cellular mechanisms of tolerance, 925–926
 - postoperative patient care protocols, 927–929
 - supraspinal mechanisms of opioids, 925
- Opioids, and the immune system, 843
- Orthopedic anesthesia, 771–931
 - fall reduction program for lower-extremity joint arthroplasty patients, **853–864**
 - assessment, 856
 - current guidelines, 854–855
 - example of selective interventions for, 858–861
 - implementation, 858
 - multidisciplinary approach, 855
 - risk factors, 856–857
 - strategies and interventions, 857
 - sustainability, 858
 - for hip fractures, **823–839**
 - challenges of, 823–824
 - institutional guidelines, 833
 - intraoperative considerations, 831–833
 - perioperative risk and optimization, 824–831
 - postoperative care, 833
 - for total joint arthroplasty, **865–880**
 - in opioid-tolerant patients, **923–932**
 - background on clinical studies, 926–927

- cellular mechanisms of tolerance, 925–926
- postoperative patient care protocols, 927–929
- supraspinal mechanisms of opioids, 925
- orthopedic trauma, **789–808**
 - advantages of regional anesthesia in, 792–793
 - benefits of regional anesthesia and analgesia in, 792
 - hemostasis in, 804
 - history of pain relief for on the battlefield, 790–791
 - local anesthetics and analgesic adjuvants, 797–798
 - neuraxial blocks, 797–798
 - peripheral nerve blocks, 798
 - neurochemistry of nociception in, 791–792
 - perioperative regional techniques in, 793–797
 - for crush injuries, 797
 - neuraxial, 793
 - paravertebral block, 793–794
 - risks of regional anesthesia in, 798–804
 - compartment syndrome, 798–800
 - in heavily sedated patients, 802–804
 - infection, 800–801
 - peripheral nerve injury, 801–802
 - role of anesthesiologist in, 789–790
- outcomes related to regional anesthesia, **809–821**
 - economic advantages, 814–815
 - improved functional outcome and inpatient fall risk, 815–816
 - length of stay, 815
 - morbidity, mortality, and complications, 811–812
 - pain management, 810–811
 - patient comfort and satisfaction, 813–814
 - resources use, transfusion, critical care admission, and ventilation, 812–813
- preoperative evaluation and patient preparation for, **881–892**
 - blood conservation strategies, 888
 - in ankylosing spondylitis, 884–885
 - in hemophilia, 886–887
 - in rheumatoid arthritis, 885–886
 - in scoliosis, 885
 - medication management, 887–888
 - patient education, 888–889
 - patients with hip (femur) fractures, 883–884
 - preoperative testing, 884
 - risk stratification, 882–883
- relationship to cancer recurrence and infection, **841–851**
 - immune system and cancer, 842
 - meta-analysis, 847
 - opioids and immune system, 843
 - regional and local anesthetic effect on immunosuppression, 843
 - studies investigating, 844–847
 - surgical site infections and regional anesthesia, 847–848
 - surgical stress leading to immunosuppression, 842–843
- setting up acute pain management service, **893–910**
 - continuous peripheral nerve blocks, 903–907

Orthopedic (*continued*)

- intravenous patient-controlled analgesia, 896–898
- multimodal approach to pain management, 894
- neuraxial analgesia, 900–903
- opioid-tolerant patients, 898–900
- role of nurses in and use of protocols, 896
- structure and individual roles within, 894–896
- setting up an ambulatory regional anesthesia program for orthopedic surgery, **911–921**
 - postoperative considerations for ambulatory perineural catheters, 918
 - potential advantages of regional anesthesia, 911–913
 - practical details for, 913–918
- ultrasound-guided regional anesthesia, **771–787**
 - central neuraxial anesthesia, 774–778
 - continuous peripheral nerve blocks, 779–780
 - evidence for patient safety and, 778–779
 - local anesthetic distribution, 780–782
 - vs peripheral nerve stimulation for peripheral nerve blocks, 773–774
- Outcomes, related to regional anesthesia for orthopedic surgery, **809–821**
 - economic advantages, 814–815
 - improved functional outcome and inpatient fall risk, 815–816
 - length of stay, 815
 - morbidity, mortality, and complications, 811–812
 - pain management, 810–811
 - patient comfort and satisfaction, 813–814
 - resources use, transfusion, critical care admission, and ventilation, 812–813

P

- Pain management, perioperative, for orthopedic surgery in opioid-tolerant patients, **923–932**
 - background on clinical studies, 926–927
 - cellular mechanisms of tolerance, 925–926
 - postoperative patient care protocols, 927–929
 - supraspinal mechanisms of opioids, 925
- setting up an acute pain service, **893–910**
 - continuous peripheral nerve blocks, 903–907
 - monitoring for falls in patients with, 906–907
 - potential complications of, 903–906
 - role in ambulatory continuous perineural catheter service, 907
 - role of local anesthetic additives in, 907
 - intravenous patient-controlled analgesia, 896–898
 - multimodal approach to pain management, 894
 - neuraxial analgesia, 900–903
 - opioid-tolerant patients, 898–900
 - role of nurses in and use of protocols, 896
 - structure and individual roles within, 894–896
 - with regional anesthesia for orthopedic surgery, 810–811
- Paravertebral block, in orthopedic trauma, 793–797
 - lower extremity, 794
 - single injection vs continuous catheter techniques, 794–797
 - upper extremity, 793–794

- Patient education, prior to orthopedic surgery, 888–889
 - in ambulatory regional anesthesia program, 913
- Patient-controlled analgesia, intravenous, in acute pain management service, 896–898
- Peripheral nerve blocks, continuous, evidence for ultrasound guidance and, 779–780
 - continuous, in an acute pain service, 903–907
 - monitoring for falls in patients with, 906–907
 - potential complications of, 903–906
 - role in ambulatory continuous perineural catheter service, 907
 - role of local anesthetic additives in, 907
 - in ambulatory anesthesia program for orthopedic surgery, **911–921**
 - ultrasound guidance vs peripheral nerve stimulation for, 773–774
- Peripheral nerve stimulation, vs ultrasound guidance for peripheral nerve blocks, 773–774
- Postoperative nausea and vomiting (PONV), decrease in with regional anesthesia, 912
- Preoperative evaluation, and patient preparation for orthopedic surgery, **881–892**
 - blood conservation strategies, 888
 - in ankylosing spondylitis, 884–885
 - in hemophilia, 886–887
 - in rheumatoid arthritis, 885–886
 - in scoliosis, 885
 - medication management, 887–888
 - patient education, 888–889
 - patients with hip (femur) fractures, 883–884
 - preoperative testing, 884
 - risk stratification, 882–883

R

- Regional anesthesia, outcomes of orthopedic surgery related to, **809–821**
 - economic advantages, 814–815
 - improved functional outcome and inpatient fall risk, 815–816
 - length of stay, 815
 - morbidity, mortality, and complications, 811–812
 - pain management, 810–811
 - patient comfort and satisfaction, 813–814
 - resources use, transfusion, critical care admission, and ventilation, 812–813
- relationship to cancer recurrence and infection, **841–851**
 - immune system and cancer, 842
 - meta-analysis, 847
 - opioids and immune system, 843
 - regional and local anesthetic effect on immunosuppression, 843
 - studies investigating, 844–847
 - surgical site infections and regional anesthesia, 847–848
 - surgical stress leading to immunosuppression, 842–843
- role in orthopedic trauma, **789–808**
 - advantages of regional anesthesia in, 792–793
 - benefits of regional anesthesia and analgesia in, 792
 - hemostasis in, 804
 - history of pain relief for on the battlefield, 790–791
 - local anesthetics and analgesic adjuvants, 797–798
 - neuraxial blocks, 797–798

Regional (*continued*)

- peripheral nerve blocks, 798
- neurochemistry of nociception in, 791–792
- perioperative regional techniques in, 793–797
 - for crush injuries, 797
 - neuraxial, 793
 - paravertebral block, 793–797
- risks of regional anesthesia in, 798–804
 - compartment syndrome, 798–800
 - in heavily sedated patients, 802–804
 - infection, 800–801
 - peripheral nerve injury, 801–802
- role of anesthesiologist in, 789–790
- setting up an ambulatory program for orthopedic surgery, **911–921**
 - postoperative considerations for ambulatory perineural catheters, 918
- potential advantages of regional anesthesia, 911–913
 - decreases postoperative nausea and vomiting, 912
 - improves patient satisfaction, 913
 - increases efficiency, 912
 - optimizes analgesia, 911–912
 - prevents unanticipated admissions and leads to earlier discharge, 912
 - surgeons preference for, 913
- practical details for, 913–918
 - equipment and personnel, 914
 - models of practice, 915–916
 - multimodal approach and clinical pathways, 916
 - patient education and screening, 913
 - postoperative conditions, 916–918
 - procedures amenable to perineural blockade, 915
 - techniques for block placement, 914–915
- ultrasound-guided, evidence for, **771–787**
 - central neuraxial anesthesia, 774–778
 - continuous peripheral nerve blocks, 779–780
 - evidence for patient safety and, 778–779
 - local anesthetic distribution, 780–782
 - vs peripheral nerve stimulation for peripheral nerve blocks, 773–774
- Rheumatoid arthritis, preoperative evaluation for orthopedic surgery in patients with, 885–886
- Risk stratification, preoperative, in orthopedic surgery patients, 882–883

S

- Safety, patient, evidence for ultrasound guidance vs perineural nerve stimulation, 778–779
- Satisfaction, patient, increase in, with regional anesthesia, 913
 - with regional anesthesia for orthopedic surgery, 813–814
- Scoliosis, preoperative evaluation for orthopedic surgery in patients with, 885
- Sedation, heavy, risk with regional anesthesia for orthopedic trauma, 802–804
- Stress response, surgical stress leading to immunosuppression via, 842–843
- Surgical site infections, and regional anesthesia, 847–848
- Surgical stress, leading to immunosuppression, 842–843

T

Total joint replacement, developing a fall reduction program for patients undergoing, **853–864**

Transfusions, with regional anesthesia for orthopedic surgery, 812–813

Trauma, orthopedic, role of regional anesthesia in, **789–808**

- advantages of regional anesthesia in, 792–793

- benefits of regional anesthesia and analgesia in, 792

- hemostasis in, 804

- history of pain relief for on the battlefield, 790–791

- local anesthetics and analgesic adjuvants, 797–798

 - neuraxial blocks, 797–798

 - peripheral nerve blocks, 798

- neurochemistry of nociception in, 791–792

- perioperative regional techniques in, 793–797

 - for crush injuries, 797

 - neuraxial, 793

 - paravertebral block, 793–797

- risks of regional anesthesia in, 798–804

 - compartment syndrome, 798–800

 - in heavily sedated patients, 802–804

 - infection, 800–801

 - peripheral nerve injury, 801–802

- role of anesthesiologist in, 789–790

U

Ultrasound-guided regional anesthesia, evidence for, **771–787**

- central neuraxial anesthesia, 774–778

- continuous peripheral nerve blocks, 779–780

- evidence for patient safety and, 778–779

- local anesthetic distribution, 780–782

- vs peripheral nerve stimulation for peripheral nerve blocks, 773–774

V

Ventilation, with regional anesthesia for orthopedic surgery, 812–813

Vomiting, postoperative, decrease in with regional anesthesia, 912